

Are graduated compression stockings still essential for VTE prophylaxis in general surgical patients?*

S. L. Schulz¹, B. Steinhilber², U. Saebler¹, S. Müller¹, D. Andresen¹, U. Boer²
(Departments of ¹Internal Medicine, ²General Surgery, Veterans Medical Center Am Urban, Berlin, Germany)

Keywords

venous thromboembolism, VTE prophylaxis, thromboelastic weight loading, graduated compression stockings, pulmonary embolism

Summary

Background: Current guidelines for the prevention of venous thromboembolism in surgical patients continue to recommend the use of graduated compression stockings (GCS) below mechanical weight loading (MMWL). However, the studies documenting a prophylactic benefit have been largely inconclusive and often of low methodological quality.
Primary methods: We studied the rate of asymptomatic venous thromboembolism (VTE) occurring in patients of a general surgery department during two consecutive five periods of 24 months with different prophylactic regimens. During the first 24 months (Group A), preventive measures consisted of MMWL and graduated compression stockings. During the second 24 months (Group B) only MMWL was used.
Results: In Group A there were 31 VTE patients with a total of 3538 operations. In Group B there were 29 VTE patients with a total of 2971 operations. 52.1% of the patients in Group A and 54.2% in Group B received VTE prophylaxis (gradings and MMWL in A, MMWL only in B). Symptomatic pulmonary embolism (PE) occurred in 45 patients in Group A and in 17 in Group B. Examination of these patients revealed a VTE in 74 and 57 patients.
Conclusions: The incidence of asymptomatic venous thromboembolism in patients of a general surgery department was high for both groups, but did not differ significantly in patients of MMWL. Graduated compression stockings did not yield additional benefits in our patients.

Schlüsselwörter

Venöse Thromboembolie, VTE Prophylaxe, mechanisches Gewichtsbelastung, Kompressionsstrümpfe, Lungenembolie

Zusammenfassung

Hintergrund: Aktuelle Leitlinien zum Thema Thromboembolieprophylaxe empfehlen auch bei nur der Fixität von Thrombozytenglykoproteinrezeptoren assoziiert zu stabilen Aktivierten Thrombin (ATP). Für alle Studien, die einen Nutzen dieser Strümpfe aufzeigen, wurden bei Erhebung der ATW durchgeführt werden. **Primäre Methoden:** Wir untersuchten die Häufigkeit von asymptomatischen Lungenembolien und Venenthrombosen bei Patienten einer Allgemeinchirurgie über zwei aufeinanderfolgende 24 Monate (Gruppe A) mittels der Thrombozytenglykoproteinrezeptoren und Kompressionsstrümpfe, während des zweiten 24 Monate (Gruppe B) ausschließlich MMWL eingesetzt. **Ergebnisse:** In Gruppe A wurden 31 VTE Patienten behandelt mit insgesamt 3538 Operationen. In Gruppe B 29 VTE Patienten mit insgesamt 2971 Operationen. 52,1% der Patienten in Gruppe A und 54,2% in Gruppe B erhielten Thrombozytenglykoproteinrezeptoren und MMWL in A, nur MMWL in B. 40 Patienten in Gruppe A und 47 in Gruppe B entwickelten Symptome einer Venenthrombose oder einer Lungenembolie. Bei den weiteren Untersuchungen fand sich in Gruppe A bei diesen Patienten ein VTE in 74 und bei Gruppe B bei diesen Patienten ein VTE in 57 Patienten. **Schlussfolgerung:** Bei einem erhöhten Risiko einer postoperativen Venenthromboembolie zeigte die Verwendung von MMWL keinen Vorteil gegenüber einer prophylaktischen Kompressionsstrümpfe oder einer mechanischen Gewichtsbelastung. Graduierte Kompressionsstrümpfe ergaben keine zusätzlichen Vorteile in unseren Patienten.

Mots clés

Maladie thromboembolique veineuse, prophylaxie, ligatures à bas poids moléculaires, les dispositifs à compression élastique, embolie pulmonaire

Résumé

Contexte: Les directives actuelles pour la prévention de la maladie thromboembolique veineuse (MTEV) en chirurgie recommandent de recommander l'usage de bas à compression élastique gradués associés à un traitement par ligatures à bas poids moléculaires (LPM), en particulier dans les études, qui ont montré un bénéfice par la compression élastique, et de faire usage uniquement des LPM seuls.
Principales méthodes: Nous avons étudié la fréquence de MTEV asymptomatique survenant dans un département de chirurgie générale au cours de 2 périodes consécutives de 24 mois avec différentes méthodes prophylactiques. Au cours des 24 premiers mois (groupe A) on a utilisé des ligatures des bas à compression élastique. Au cours des 24 mois suivants (groupe B) seuls des LPM ont été utilisés.
Résultats: Chez le groupe A 31 VTE patients ont été traités au total de 3538 opérations. Chez le groupe B 29 VTE patients ont été traités au total de 2971 opérations. 52,1% des patients du groupe A et 54,2% du groupe B ont reçu une prophylaxie de la VTE. Malgré tout, 40 patients ont présenté une symptomatologie suggestive une MTE dans le groupe A et 47 dans le groupe B. L'examen a révélé 7 cas de MTE dans le groupe A et 5 dans le groupe B.
Conclusions: Le risque de la MTE élevés dans un département de chirurgie a été enregistré dans nos patients et l'utilisation de prophylaxie élastique n'a pas eu de bénéfice supplémentaire par rapport à l'utilisation de LPM. Les bas à compression élastique gradués ne semblent pas avoir apporté de bénéfice supplémentaire chez nos patients.

Are bas à compression élastique gradués still essential for the prophylaxis of the venous thromboembolism in the general surgery?

Phlébologie 2008, 37: 372-377

Schlüsselwörter zur Thrombozytenglykoproteinrezeptoren in der Allgemeinchirurgie nach publiziert?

* Diese Arbeit wurde unterstützt durch die Förderung der Deutschen Gesellschaft für Phlebologie in Mainz am 15. April 2008. Correspondence: Schulz S. L. (s.schulz@vmed.uni-berlin.de)

Received: November 19, 2007; accepted in revised form: November 21, 2007

Venous thromboembolism (VTE) continues to account for considerable morbidity, mortality and medical care expense. In the USA the incidence of deep venous thrombosis (DVT) is reported to be approximately 450 000 cases per year and the overall incidence of pulmonary embolism (PE) to be about 355 000 cases per year (8). Numerous studies have indicated that especially surgical and trauma patients are at significant risk for developing venous thromboembolic complications, including pulmonary embolism. Without prophylactic treatment, the risk of VTE (2, 19, 50) is considered to be:

- 25–30% in general surgery (3, 13, 46),
- 60% after major trauma (22, 23) and
- up to 45–70% with hip or knee replacements.

Both physical and pharmacological measures aimed at preventing thromboembolic events have been previously evaluated. In earlier studies graduated compression stockings were found to substantially reduce the incidence of VTE in surgical patients (2, 13, 30, 46, 60). During recent years, however, notable developments in pharmacological thromboprophylaxis have been achieved. The introduction of low molecular weight heparins (LMWH), in particular, has resulted in significant advances in treatment modalities so that today they have become the standard for VTE prophylaxis in most countries. The prophylactic properties of LMWHs have yielded equal or even slightly better outcomes than unfractionated heparins. Moreover, they have fewer side effects, do not require laboratory monitoring, and possess pharmacokinetic characteristics which permit their use as a subcutaneously administered, fixed daily dose. LMWHs are reported to reduce the overall incidence of deep vein thrombosis after general surgery by up to 70% (2, 33).

In light of this progress, reevaluation of the common use of graduated compression stockings, as it is still recommended in this setting, is warranted.

Patients, methods

Study population and design

The study population consisted of all patients treated in a general surgery department over a period of four years.

With a prospective observational study design, the incidence of symptomatic deep venous thrombosis and pulmonary embolism was investigated with two different VTE prophylaxis regimens. During the

- first 24 months (group A) patients at risk received combined treatment for the prevention of thromboembolism consisting of LMWH and graduated compression stockings,
- second 24 month period (group B) only LMWH was used for prophylaxis.

Treatment

Prophylactic treatment was given to all patients undergoing major surgery (>90 min). The treatment was initiated the day before surgery and was continued for five days thereafter, or until complete mobilization was achieved. The prophylactic measures were also taken in patients with minor surgery and patients who were not actually operated on if they were expected to be immobile for more than 6 hours during the day for more than two days. Patients under the age of 14 did not receive VTE prophylaxis.

The thigh-length compression stockings (TED[®] Kendall) in group A were fitted to each patient's legs according to the manufacturer's instructions. The patients were advised to wear them 24 hours a day. Compliance with the wearing of stockings was checked at least twice daily. Patients found without stockings or not wearing them properly on more than two occasions were categorized as non-compliant. Patients with known significant peripheral artery disease (intermittent claudication and/or ABI < 0.7) did not receive stockings.

The LMWH employed in both groups was enoxaparin. It was administered according to a simple weight-adjusted protocol. Patients with a body mass index (BMI)

of up to 30 kg/m² received 20 mg of enoxaparin and patients with BMI > 30 kg/m² received 40 mg – subcutaneously, once daily. Patients in whom the use of LMWH was contraindicated (those with a history of or suspected active HIT II or allergic skin reactions) were treated with 15 mg of fondaparinux subcutaneously, twice daily.

In group B, no stockings were used. The prophylactic treatment consisted of LMWH (or fondaparinux alone, administered following the same protocol as in group A).

All patients in groups A and B were strongly encouraged to get out of bed and walk as early as possible.

Outcomes

In all patients with clinical symptoms suggesting deep vein thrombosis colour-Doppler ultrasonography of the pelvic and leg veins was done, including compression testing of the leg veins. The leg examination included all veins from the thigh down to the ankle. All sonographic examinations were carried out by two highly-skilled examiners with an ATL / Philips HDI 3000 machine, using a 3–7 mHz curved array scanner for the pelvic vein and a 5–7 mHz phased array scanner for the leg veins. If this examination was temporarily not possible, or if the results were inconclusive, ascending phlebography followed.

Patients with symptoms of pulmonary embolism underwent spiral computed tomography of the lungs. This was judged positive if it revealed distinct filling defects in large vessels. If the diagnosis remained inconclusive after this examination, a ventilation-perfusion lung scan (judged positive only in cases of „high probability“ results) and/or a pulmonary angiography was performed.

Results

A total of 6187 patients (Tab. I) were treated in the general surgery department during the study period, with a total of 5861 operations.

Treatment group characteristics

In group A (treated with stockings and LMWH), there were 3181 patients: 1742 men and 1439 women, with a mean age of 48 years. The mean duration of hospital stay was 7.3 days. 2720 of the A patients had surgery with a total of 3890 operations (Tab. 1, Tab. 2). Four hundred sixty-one patients had no surgery. VTE prophylaxis (Tab. 3) was applied to a total of 2623 (82.5%) of the 3181 patients of period A: to 2422 (89.1%) patients with surgery and to 202 (43.8%) patients without surgery. Fifteen patients in group A could not wear compression stockings due to advanced occlusive artery disease and in others because of extreme obesity. Twenty-eight patients were non-compliant with wearing stockings. Three patients in group A needed to be treated with heparin, instead of LMWH, due to contraindications.

In group B (only LMWH), 2986 patients were treated: 1691 men and 1295 women, with a mean age 49 years. The mean duration of hospital stay was 6.2 days and 2551 patients had a total of 2914 operations. Four hundred thirty-three patients had no surgery. VTE prophylaxis was provided to 2512 (84.0%) of the 2986 patients in period B: to 2305 (90.2%) patients with surgery and to 207 (46.0%) patients without surgery. Two patients had to be treated with heparin, instead of LMWH.

Tab. 1
Atrial characteristics,
type of surgery

		group A	group B
		LMWH + stocking	LMWH only
patients	total number, sex	3181	2986
	men	1742	1691
	women	1439	1295
	age (years)	48	49
	days of stay	7.3	6.2
type of surgery	vascular	261	157
	thyroid	149	104
	choleci	119	62
	total colectom	49	65
	gastrointestinal	227	153
	urologic sup	99	164
	urologic	176	109
	gyn	215	205
	law	3	4
	gallbladder resected	44	29
	gallbladder sup	296	284
	pancreas	3	3
	hepatocyst resected	257	81
	hepatocyst sup	28	44
	colostomy resected	42	33
	urothelium	190	93
	cholec	226	202
	retroperitoneal lymph nodect	45	29
	duodenal resected	81	81
	stomach resected	67	173
urothelium	25	29	
total	3890	2914	

Outcomes (Tab. 4)

Deep vein thrombosis

Thirty-eight patients in group A and 40 in group B developed clinical symptoms suggestive of a deep vein thrombosis which required further investigation. Thirty patients in A and 31 patients in B received color-duplex assisted compression sonography and eight patients in A and nine in B received phlebography. A DVT was confirmed in six patients in group A and in three patients in group B.

Tab. 2
Number of operations
per patient

number of operations	group A	group B	p value
	LMWH + stocking	LMWH only	
1	2517	2349	0.658
2	127	124	0.718
3	44	45	0.658
4	18	19	0.596
5	9	9	0.893
6	3	10	0.157
7	0	1	0.202
≥8	0	1	0.202

376
Schick et al.

	VTE prophylaxis	group A	group B	P value
		LMWH + stockings	LMWH only	
patients		7181	7166	
with VTE	—	1735	2553	0.002
without VTE	+	2429 (33.8%)	2104 (29.2%)	0.01
without VTE	—	465	452	0.997
without VTE	+	2162 (30.2%)	2152 (30.1%)	0.395
total	+	2627 (36.6%)	2107 (29.4%)	0.011

—: not defined in patients.

Table 3
VTE prophylaxis

	group A	group B	P value
	LMWH + stockings	LMWH only	
operative	663 patients	663 patients	
days	3075	3117	
phlebography	671	391	
post-D	511	412	
next post-op	118	169	
post-angiography	319	171	
total patients	4677	4704	0.985
patients with VTE	7	5	0.667
incidence	1.52	1.07	0.127

Table 4
Incidence of VTE rates

Pulmonary embolism

A pulmonary embolism was suspected in six patients in group A and seven in group B during their hospital stay. 12 of these patients underwent spiral computed tomography of the lungs, one patient had a ventilation-perfusion scan and one needed an additional pulmonary angiogram. The pulmonary embolism was confirmed in one patient in group A and in three in group B.

Characteristics of patients with VTE

All the 12 patients (7 in group A, 5 in group B) with confirmed VTE were receiving prophylaxis at the time the VTE event occurred. The mean duration of hospital stay in these patients was 38 days in group A and 39 days in group B. All patients with PE in both groups had multigrain disease.

Discussion

This study, which took place over a period of four years and involved 6167 patients on a general surgery ward, revealed no significant differences in the incidence of asymptomatic venous thromboembolism whether graduated compression stockings were used in addition to LMWH prophylaxis, or not.

A review of the literature over the last three decades shows that, despite significant advancements in the prevention of venous thromboembolism, there is still a considerable degree of variation in regard to risk group definition, the number of patients receiving prophylaxis, and prophylactic modalities.

To achieve better standardization in VTE prophylaxis, several consensus conferences have generated guidelines which are aimed at promoting and attaining adequate VTE prophylaxis for the individual patient. Yet, even the latest guidelines concerning the

prevention of thromboembolism in surgical patients, namely the updated ACCP guidelines from 2004 (24) and those of the International Consensus Statement dating from 2006 (46), are not completely concordant. In regard to general surgery patients, the former do not recommend a specific prophylaxis in low-risk patients, propose the alternative use of stockings or heparin in moderate- and high-risk patients, and recommend combination therapy only in very high-risk patients. The latter consider the use of compression stockings in low risk patients, recommend stockings with intermittent pneumatic compression as an alternative to LMWH in moderate risk patients with bleeding tendency and only consider compression stockings as an adjunct to LMWH in high-risk patients. In other fields of surgery (for example oncologic surgery or gynecologic surgery) there is even less evidence for the necessity of compression stockings. In these fields the recommendations are either extrapolated from general surgery or consensus concerning the use of compression stockings are lacking - as it also the case in all recently published studies on VTE prophylaxis in medical patients (26, 41, 52).

There are generally nine studies cited which substantiate significant VTE risk reduction following the use of graduated compression stockings in general surgery (65). All of them were carried out 10 or more years ago (3, 4, 28, 43, 45, 46, 58, 63, 64). The study designs are rather heterogeneous and most of them included a total of less than 100 patients (5, 28, 45, 46, 58). Only two out of the nine studies (3, 28) compared the use of stockings with the absence of prophylactic measures. These found a 50% reduction in the rate of venous thromboembolism in patients who wore stockings. In the seven other studies, a combination therapy was used consisting of either intermittent pneumatic compression (43, 58) or pharmacological methods (dextran, low dose heparin, dilydrogesterone-heparin) together with graduated compression stockings (3, 43, 45, 63, 64). In five of the studies the effectiveness of compression stockings was investigated on only one of the patient's legs; the other leg served as a control (5, 43, 55, 56, 58).

Several studies in the following years have provided evidence that low dose heparin significantly reduced the incidence of postoperative VTE, and the benefit appeared to increase with the additional use of compression stockings (11, 15, 31, 32, 45). Subsequently, low dose heparin in combination with compressive stockings became the most widely adopted measure for VTE prophylaxis in general surgery – a regimen which has been used in many facilities until today (4, 10, 12, 65).

When LMWHs were introduced into VTE prophylaxis in the mid-1990s, they first proved their effectiveness in orthopedic surgery, namely with hip and knee replacements. Later LMWHs were adopted in general surgery and were combined with the compression stocking prophylaxis already in use. Only four studies can be found which show an advantage to the combination of LMWH with compression stockings over stockings alone – none of them in general surgery (1, 38, 42, 47). In all four studies the group employing compression stockings alone is categorized as a placebo group. However, the incidence of VTE in these placebo groups is actually about the same as in previous studies where no prophylactic measures were used. Only one single, rather small study has compared, as we have, the use of LMWH in combination with compression stockings to LMWH alone for VTE prophylaxis. Katsifiki et al. (34), randomized 78 patients undergoing a total hip replacement into three groups:

- a control group without prophylaxis,
- a group which received enoxaparin once daily,
- a group which received enoxaparin and wore graduated compression stockings (TED) for a period of 8–10 days.

All patients had a preoperative perfusion leg scan and a postoperative perfusion-ventilation scan, as well as bilateral ascending phlebography between days 8 and 12. The DVT rates were 93%, 38% and 25%, respectively. No other study comparing LMWH with and without stockings has been published as of today – neither in general surgery nor in any other medical specialty.

The mechanism of action through which compression stockings prevent thrombo-

embolism remains unclear, though it is thought to be multi-functional with a reduction in venous diameter playing the key role. External compression was shown to reduce the overall cross-sectional area of the lower limb and increase the flow velocity within the veins. Compression also appears to improve the evacuation and reabsorption of incompetent and incompletely emptied valvular cusps. Reduction in reflux-related stasis and augmented flow velocity could decrease the risk of thrombus formation by reducing venous wall distension, local contact time, and the concentration of coagulation reactants (2, 14, 57, 61, 66).

Despite their widespread use, there is also no agreement as to the most effective stocking design to achieve the above mentioned effects. High-length TED stockings, which are probably employed in most studies, have intended pressures of 18 mmHg at the ankle, 14 mmHg at the calf, 8 mmHg at the knee, 0 mmHg above the knee, and 0 mmHg at the thigh. It is not known whether these pressures are optimal. Moreover, some authors estimate calf-length stockings to be more effective than thigh-length and some prefer non-elastic ones (7, 9, 26). Stocking design is not always specified in the study protocols and, particularly in some of the newer studies, it is not clear whether or not stockings were used at all (8, 27, 39, 44, 59, 41, 35, 32).

Even those stockings believed to be ideally designed may not achieve the warranted effects. A study measuring pressure gradients in patients wearing prophylaxis stockings showed that 90% of the stockings failed to produce the ideal pressure gradient of 18–14–8 from the ankle to the knee; 54% even produced an inverse gradient (7, 67).

Given the recommended pressure gradients intended to be realized by the design of prophylaxis stockings, it remains obscure why other types of compression stockings, e.g., those recommended for the prevention of post-thrombotic syndrome following a deep vein thrombosis, achieve significantly higher pressure ranges (mostly 40–30–21 mmHg) although they must be worn for much longer time periods. These higher pressure compressive stockings fail, however, to prevent DVT recurrence, whereas

prophylaxis stockings are claimed to reduce DVT incidence by 50% or more (9, 37).

Finally, prophylaxis stockings were considered for supine patients (61). However, today most patients are mobilized within only a few hours after surgery. Nevertheless guidelines recommend stockings for 5–7 days (61).

Unlike most other studies, ours did not serially screen all our study patients for deep venous thrombosis or pulmonary embolism. The significance of an asymptomatic VTE detected by screening methods such as ^{99m}Tc fibrinogen marking, phlebimetry, contrast phlebography, compression sonography, D-dimer testing, or ventilation-perfusion scans is a matter of debate (6, 35, 40, 54). Conclusive evidence that treatment of asymptomatic patients with VTE detected by these methods results in better outcomes is lacking, not only in regard to nonfatal or fatal pulmonary embolism (35), but also in regard to postthrombotic syndrome (25) and pulmonary hypertension (48). Recent study designs have begun to focus more, as we have, on symptomatic thromboembolic events as clinically relevant endpoints (41).

Limitations

We cannot rule out the possibility of diagnostic bias in our study, since the use of stockings cannot be blinded and for organizational reasons we were not able to perform randomization. It is possible that physicians were more likely to order an imaging test for patients who had not received compression stockings than for those who had. Nevertheless, the incidence of asymptomatic VTE in our surgical patients was very low with both prophylaxis regimens.

A decline in VTE rates in clinical medicine has been anticipated in recent years (17, 23). Research in this area is lacking, but considering the still doubtless significance of the Virchow triad in regard to VTE pathogenesis, the current prevailing concept of early mobilization is likely to be the most effective measure contributing towards the prevention of thromboembolism in all areas of medicine. New surgical techniques, especially endoscopic and fast track surgery and modern anesthesiologic methods, are

also reported to reduce VTE rates (49, 51, 53). Furthermore, complicated risk stratification concepts are increasingly being disregarded in favor of simple and effective protocols like the one we used with our surgical patients (24) and finally, new, even more potent, anti-thrombotic agents are being developed (16, 28, 21, 49).

Conclusion

Our data showed that the rate of symptomatic venous thromboembolism in patients in a general surgery ward is kept low when a concept of early mobilization and a simple, but strictly applied protocol of preventative low molecular weight heparin is implemented. Graduated compression stockings did not yield an additional preventive effect with this regimen.

References

1. Agelli G, Perrella F, Biondinianni P, Sestini P, Pini M, P. Angeli A, et al. Enoxiparin plus compression stockings compared with compression stockings alone in the prevention of venous thromboembolism after elective nonorthopedic surgery. *N Engl J Med* 1999; 340: 86-92.
2. Agui G, Harthorn G, Baker D. Graduated compression stockings in the prevention of venous thromboembolism. *Br J Surg* 1999; 86: 1002-1004.
3. Altai A, Williams D, Butler D, Le Quang L P. The use of graduated compression stockings in the prevention of postoperative deep vein thrombosis. *Br J Surg* 1995; 82: 172-174.
4. Arangis SS, Lee TA. Elastic compression stockings for prevention of deep vein thrombosis. *Cochrane Database Syst Rev* 2003; 3: CD001484.
5. Duggan B, Lindblad B. The thromboprophylactic effect of graded compression stockings in combination with Acton. *Br Arch Surg* 1994; 119: 1329-1331.
6. Brennan R, Bain S, Brumby R, Sang G, Dore S, G. Cattan D. Venous sonography for the diagnosis of asymptomatic venous thrombosis in patients with cancer undergoing chemotherapy. *J Ultrasound Med* 2004; 23: 675-678.
7. Borel A, Williams S, Cramer A, Shaw B, Strong PJ, Birk AC. Elastic compression stockings in elective orthopedic surgery. An assessment of the in vivo performance of commercially available stockings in patients having hip and knee arthroplasty. *J Bone Joint Surg Br* 2000; 82: 116-119.
8. Birk AC, Kaplan EL. Thromboprophylaxis in surgical patients. *Per J Med Res* 2004; 9: 104-111.

9. Dimarco B, Butler DR, Harvey H, O'Rourke MV, de Bak M. Leg Band. Randomized trial of effect of compression stockings in patients with symptomatic proximal vein thrombosis. *Lancet* 1997; 349: 790-792.
10. Caputo D, Anselmi J, Sghele LR, Colan FB, Rossi D. The use of low molecular weight heparin for the prevention of postoperative venous thromboembolism in general surgery: A survey of practice in the United States. *Br J Surg* 2002; 89: 58-60.
11. Clagett GP, Rasch R. Prevention of venous thromboembolism in general surgical patients. Results of meta-analysis. *Ann Surg* 1998; 228: 227-240.
12. Clagett GP. Prevention of postoperative venous thromboembolism: An update. *Am J Surg* 1998; 168: 745-752.
13. Collins GA, Taylor RL, Ortiz G. Rates of venous thrombosis after general surgery. Combined results of randomized clinical trials. *Lancet* 1996; 348: 143-146.
14. Colletto Smith PH, Harty D, Scurr D. Deep vein thrombosis: effect of graduated compression stockings on duration of deep vein of the calf. *Br J Surg* 1994; 78: 724-726.
15. Collins R, Scraggs A, Yusuf S, Peto R. Reduction of fatal pulmonary embolism and venous thrombosis by preoperative administration of subcutaneous heparin. Overview of clinical randomized studies in general orthopedic and oncologic surgery. *N Engl J Med* 1998; 338: 1162-1173.
16. Colwell CW, Berkowitz SD, Davidson BL, Lello PA, Ginsberg JS, Lefkowitz DR, et al. Comparison of Ximelupatan as ambulatory thrombolytic effect with enoxaparin for the prevention of venous thromboembolism following total hip replacement. A randomized double-blind study. *J Thromb Haemost* 2004; 3: 2119-2126.
17. Denti R, Maney Jone B. Incidence of thromboembolism after transarterial resection of the prostate (TURP) - a study on TED stocking prophylaxis and haemostatic review. *Scand J Urol Nephrol* 2002; 36: 119-121.
18. Durrant H, Lindbloom PW, Quinlan DR, Wilton AN, Gougher MN. Short-duration prophylaxis against venous thromboembolism after total hip or knee replacement. A meta-analysis of prospective studies investigating symptomatic outcomes. *Arch Intern Med* 2002; 162: 1465-1471.
19. Eichinger S, Kralj PA. Prevention of deep vein thrombosis in orthopedic surgery. *Per J Med Res* 2004; 9: 112-118.
20. Eriksson BI, Bauer KA, Lassen MR, Bajic AI. Enoxaparin compared with enoxaparin for the prevention of venous thromboembolism after hip fracture surgery. *N Engl J Med* 2001; 345: 1296-1304.
21. Evans RC, Price CM, Keele B. Ximelupatan/Ximelupatan a review of its use in the prevention of venous thromboembolism in orthopedic surgery. *Thromb* 2004; 64: 640-670.
22. Green WH, Ay RM, Cole RL, Chan F, Szabo DF, Sobel EA, et al. A comparison of low-dose heparin with low molecular weight heparin or prophylax

- is against venous thromboembolism after major surgery. *N Engl J Med* 1996; 335: 761-767.
23. Green WH, Chan F, Clagett GP. Prevention of venous thromboembolism. *Chest* 2001; 118: 1328.
24. Green WH, Pinedo GC, Bar JA, Bergqvist D, Lassen MR, Colwell CW, et al. Prevention of venous thromboembolism: The Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126: 376a-400a.
25. Ginsberg JS, Tackles J, Butler DR, Mac Sween R, Singer S, Durrant H. Post-thrombotic syndrome after hip and knee arthroplasty: a cross-sectional study. *Arch Intern Med* 2000; 160: 699-702.
26. Harwood MJ, Lawson DL, Tomchasz EJ, Goldberg PJ. Short knee-length versus long-length graduated compression stockings in the prevention of deep venous thrombosis. *Semin Surg* 2002; 30: 15-16.
27. Hillman M, Lata L, Sotomoni K, Lehmann J, Nagel S, Korte M. Enoxaparin vs heparin for prevention of deep vein thrombosis in acute ischemic stroke: a randomized double-blind study. *Ann Neurol* 2002; 51: 94-97.
28. Holbrook CJ. Graduated compression for preventing deep-vein thrombosis. *Drugs* 1996; 52: 969-976.
29. Horlander KJ, Mezzo DM, Lopez CV. Pulmonary embolism mortality in the United States: 1979-1998: an analysis using multiple cause mortality data. *Arch Intern Med* 2003; 163: 1711-1717.
30. Jeffrey PC, Nicholas AN. Graduated compression stockings in the prevention of postoperative deep vein thrombosis. *Br J Surg* 1998; 77: 340-343.
31. Kakkar VV, Connors T, Speroff T, Brooks DR, Hirsh J, Colwell CW, Wessler S, Yu ET. Efficacy of low doses of heparin in prevention of deep vein thrombosis after major surgery. A double-blind randomized trial. *Lancet* 1972; 2: 391-396.
32. Kakkar VV, Connors T, Brooks DR, Bullard J, Therasil P. Prevention of deep venous thrombosis by low doses of heparin. Reappraisal of results of international multicentric trial. *Lancet* 1977; 1: 567-569.
33. Kakkar VV, De Lorenzo F. Prevention of venous thromboembolism in general surgery. *Baillieres Clin Haematol* 1998; 11: 605-619.
34. Kalko H, Koppertgen BA, Nicholas AN, Garcia J, Hill S, Lopez F, et al. Deep vein thrombosis prophylaxis with low molecular weight heparin and elastic compression stocking for total hip replacement. A randomized controlled trial. *Int J Angiol* 1996; 55: 161-169.
35. Keenan C. Noninvasive diagnosis of deep vein thrombosis in postoperative patients. *Semin Thromb Haemost* 2004; 31: 3-9.
36. Kiefer PA, Witt C, Vogel G, Koppertgen R, Schomker U, Therasil W. A randomized comparison of enoxaparin with other anticoagulants for the prevention of venous thromboembolism in medical patients with heart failure or severe respiratory disease. The Prince study group. *Am Heart J* 2003; 145: 624-631.
37. Keele B, Scahill MW, Houshyk R, Narayan SA, Price CM. Non-pharmacological measures for prevention of post-thrombotic syndrome. *Cochrane Database Syst Rev* 2004; 1: CD004174.